

# A Population Study: HLA-A, -B, -C, -DRB1, -DQB1, -DQA1 Alleles, and ABDR Haplotype Frequencies in the Aegean Region of Türkiye

## Bir Popülasyon Çalışması: Türkiye'de Ege Bölgesi'nde HLA-A, -B, -C, -DRB1, -DQB1, -DQA1 Alleles ve ABDR Haplotiplerinin Frekansı

Aslı ÖZKIZILCIK KOÇYİĞİT<sup>1</sup>, Tülay KILIÇASLAN AYNA<sup>1</sup>, Mustafa SOYÖZ<sup>1</sup>, Burcu ÇERÇİ ALKAÇ<sup>1</sup>, Rasime Derya GÜLEÇ<sup>2</sup>, Deniz İlhan TOPÇU<sup>3</sup>, İbrahim PİRİM<sup>1</sup>

<sup>1</sup>İzmir Katip Çelebi University Faculty of Medicine, Department of Medical Biology, İzmir, Türkiye

<sup>2</sup>İzmir City Hospital, Tissue Typing Laboratory, İzmir, Türkiye

<sup>3</sup>İzmir City Hospital, Clinic of Medical Biochemistry, İzmir, Türkiye

**Cite as:** Özkızılcık Koçyigit A, Kılıçaslan Ayna T, Soyöz M, et al. A population study: HLA-A, -B, -C, -DRB1, -DQB1, -DQA1 alleles, and ABDR haplotype frequencies in the Aegean Region of Türkiye. Forbes J Med. 2025;6(1):1-8

### ABSTRACT

**Objective:** Human leukocyte anti-gen (HLA) molecules are vital molecules because they present foreign anti-gens to immune cells and trigger the immune response. Thus, they are associated with diseases, and they play crucial roles in solid organ transplantation and hematopoietic stem cell transplantation. This study aimed to determine the HLA allele and three-locus haplotype frequencies of the Turkish population. This is the first study in Türkiye to include such a large, sizable population.

**Methods:** The study included 6,039 bone marrow donors. Sequence-specific oligonucleotide probes and sequence-specific primers were employed for HLA typing. After excluding related individuals, 1222 A, 1229 B, 930 DRB1, 102 C, 227 DQB1, 163 DQA1 allele couples, and 336 ABDR haplotype couples were analyzed using Arlequin v3.5 Software and the R package programs. The results were compared with those from the border countries.

**Results:** The most frequently observed HLA alleles were A\*02, B\*35, C\*07, DRB1\*11, DQA1\*01, and DQB1\*03. The most prevalent ABDRB1 haplotype was A\*24 B\*35 DRB1\*11 in our population. For all loci, the population was in Hardy-Weinberg equilibrium. A linkage disequilibrium analysis revealed a strong link between the A-B, A-DRB1, and B-DRB1 loci. The results of the two methods were similar.

**Conclusion:** These findings are consistent with populations in bordering nations. Our findings will support numerous therapeutic applications, including drug discovery, disease therapy, and organ transplantation. The Arlequin and R programs are suitable for analyses.

**Keywords:** HLA, allele frequency, haplotype frequency

Received/Geliş: 10.10.2024

Accepted/Kabul: 20.12.2024

Epub: 14.02.2025

Publication Date/  
Yayınlanma Tarihi: 10.04.2025

Corresponding Author/  
Sorumlu Yazar:

Aslı ÖZKIZILCIK KOÇYİĞİT, MD,

İzmir Katip Çelebi University  
Faculty of Medicine, Department of  
Medical Biology, İzmir, Türkiye

✉ aslikocyyigit2021@gmail.com

ORCID: 0000-0002-0015-7070

### ÖZ

**Amaç:** İnsan lökosit antijeni (HLA) molekülleri yabancı antijenleri immün hücrelere sundukları ve immün yanıtı başlatmaları açısından önemli moleküllerdir. Bu sebeple, hastalıklarla ilişkili oldukları gibi solid organ ve hematopoetik hücre nakillerinde kritik rol oynarlar. Bu çalışmada, Türk popülasyonundaki HLA allel ve 3 lokus haplotip frekanslarının belirlenmesi amaçlanmıştır. Bu çalışma Türkiye'de yapılmış en büyük popülasyonu içermektedir.

**Yöntem:** Bu çalışmaya 6.039 kemik iliği vericisi dahil edilmiştir. HLA tiplemesi sekans spesifik oligonükleotit prob ve sekans spesifik primer yöntemleriyle gerçekleştirilmiştir. Akriba kişiler hariç tutulduktan sonra toplamda 1222 A, 1229 B, 930 DRB1, 102 C, 227 DQB1 ve 163 DQA1 allel çiftiyle birlikte 336 ABDR haplotip çifti Arlequin v3.5 Programı ve R paket program ile analiz edilmiştir. Sonuçlar komşu ülkelerdeki frekanslarla karşılaştırılmıştır.



Copyright© 2025 The Author. Published by Galenos Publishing House on behalf of Buca Seyfi Demirsoy Training and Research Hospital. This is an open access article under the Creative Commons AttributionNonCommercial 4.0 International (CC BY-NC 4.0) License.

Copyright© 2025 Yazar. Buca Seyfi Demirsoy Eğitim ve Araştırma Hastanesi adına Galenos Yayınevi tarafından yayımlanmıştır. Creative Commons Atıf-GayriTicari 4.0 Uluslararası (CC BY-NC 4.0) Uluslararası Lisansı ile lisanslanmış, açık erişimli bir makedir.



**Bulgular:** Popülasyonumuzda en sık saptanan HLA alleleri A\*02, B\*35, C\*07, DRB1\*11, DQA1\*01 ve DQB1\*03 olarak bulundu. En sık prevalansa sahip ABDRB1 haplotipi ise A\*24 B\*35 DRB1\*11 olarak belirlendi. Bütün lokuslar için popülasyonumuz Hardy-Weinberg eşitliğine uymaktaydı. Bağlantı eşitsizliği analizine göre A-B, A-DRB1 ve B-DRB1 lokusları arasında güçlü bir bağlantı olduğunu gösterdi. Her iki istatistiksel analiz sonucu da aynı çıktı.

**Sonuç:** Bulgular komşu ülkelerle benzer oranda çıkmıştır. Bu bulgular ilaç keşfi, hastalık tedavisi ve organ nakli gibi birçok terapötik uygulamayı destekleyici niteliktedir. Bu istatistiksel analizler için Arlequin ve R programlarının uygun olduğu görülmüştür.

**Anahtar Kelimeler:** HLA, allel frekansı, haplotip frekansı

## INTRODUCTION

The *human leukocyte anti-gen (HLA)* gene on chromosome 6p21 encodes HLA molecules, which are cell surface proteins that are crucial for anti-gen presentation to T lymphocytes.<sup>1</sup> HLA gene Regions are the most polymorphic on the human genome, and HLA molecules expressed from these Regions play critical roles in the immune system.<sup>2</sup> HLA is divided into two classes: HLA Class 1 (HLA-A, HLA-B, and HLA-C) and HLA Class 2 (HLA-DR, HLA-DQ, and HLA-DP).<sup>3</sup> Type 1 HLA molecules present the anti-genic peptides to CD8+ T-cells, whereas type 2 HLA molecules present the anti-genic peptides to CD4+ T-cells.<sup>4</sup>

HLA typing is commonly used in clinics for hematopoietic stem cell and solid organ transplantation. Furthermore, because of its roles in immune response, it is linked to a variety of diseases. Thus, community-based studies on HLA allele frequency may be valuable for detecting genetic variation and population origin, identifying vaccine-safe epitopes, and discovering disease associations.<sup>5</sup> These studies also aid in the creation of HLA databases. The frequencies of alleles and haplotypes in various populations have been the subject of numerous studies. Research conducted both nationally and internationally has linked specific alleles to distinct clinical symptoms.<sup>6</sup>

The Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium (LD) are essential concepts in population genetics that explain genetic diversity and inheritance patterns. HWE establishes a standard for forecasting genotype frequencies in a static population, whereas LD assesses the non-random correlation of alleles across several loci. Collectively, they provide insights into genetic composition and evolutionary mechanisms.<sup>7</sup> Deviations from HWE may signify genotyping inaccuracies or the existence of uncommon variations, especially within extensive genomic datasets.<sup>8</sup> LD quantifies the extent to which alleles at distinct loci are co-inherited, influenced by factors including population structure and selection. The precise measurement of LDs is essential for applications such as disease association studies, and novel approaches have enhanced LD estimation under HWE conditions. The correlation between HWE and LD was significant, as variations from HWE might influence LD assessments and hence confound genetic investigations.<sup>9</sup>

In this study, we investigated the frequencies of alleles and haplotypes in bone marrow (BM) donors. We compared the most frequently observed alleles and haplotypes with those from neighboring nations. Furthermore, we evaluated two statistical tools (Arlequin and R) to analyze the data.

## METHODS

### Samples

Molecular HLA typing data from 6039 healthy Turkish people were retrospectively analyzed. Between 2009 and 2018, these individuals were BM donors who applied to the tissue typing laboratory at the hospital. Related individuals were excluded to ensure that genetic variation was assessed objectively and in a population-representative manner. In total, 1222 A, 1229 B, 930 DRB1, 102 C, 227 DQB1, 163 DQA1 allele couples, and 336 ABDR haplotype couples were analyzed. The results were compared with those from bordering countries.

Informed consent was provided by all patients. In accordance with the Declaration of Helsinki, our İzmir Katip Çelebi University Non-interventional Clinical Research Ethics Committee approved the study (decision no: 0097, date: 23.03.2023).

### Molecular Human Leukocyte Anti-gen Typing

Sequence-specific oligonucleotide probe (SSOP) and sequence-specific primer (SSP) methods were used for the molecular typing of HLA. Deoxyribonucleic acid (DNA) was isolated using an EZ1 DNA Blood Kit (Qiagen, Hilden, Germany) via an automatic system (Geno M-6, Qiagen). Commercial SSOP (Lifecodes, Rodermark, Germany) and SSP kits (Olerup, Stockholm, Sweden) were used for the procedures. The tests were performed according to the manufacturer's instructions. Briefly, the Regions were amplified by polymerase chain reaction (PCR), and the amplicons were hybridized with specific probes for the SSOP method. The results were analyzed using a Luminex fluoroanalyzer. For the SSP method, samples were amplified using SSPs by PCR, and the amplicons were evaluated on agarose gels using score software.

### Statistical Analysis

All statistical analyses, including total allele frequencies, haplotype frequencies LD, and HWE tests, were conducted

using R 4.2 programming language with the pegas and adagen packages, and Arlequin v3.5.<sup>10-13</sup>

R is a robust and versatile programming language and software environment, specifically designed for statistical analysis, data science, and visualization. Arlequin, on the other hand, is a software specifically designed for the examination and interpretation of genetic data. The tool provides a diverse range of statistical methodologies and tests to facilitate the comprehension of genetic population structure, genetic diversity, and demographic history. The congruent outcomes of these two systems are of significant importance for their application, particularly in the analysis of HLA allele and haplotype frequencies. This comparison enables the assessment of the trade-offs between a versatile, general-purpose tool like R and a specialized software platform for population genetics such as Arlequin. This facilitated the identification of the most efficient, precise, and user-centric method for evaluating HLA allele and haplotype frequencies in our study.

## RESULTS

In total, 19 different HLA-A, 28 different HLA-B, 14 different HLA-C, 13 different HLA-DRB1, 6 different HLA-DQA1, and 5 different HLA-DQB1 alleles were identified. The allele with the highest frequency was HLA-A\*02 [allele frequency (AF): 0.195990], followed by HLA-A\*24, HLA-A\*03, HLA-A\*01, and HLA-A\*11 (AF: 0.147709, 0.126432, 0.119885, and 0.087561), respectively. The five initial HLA-B alleles that were most frequently found were HLA-B\*35, HLA-B\*51, HLA-B\*44, HLA-B\*18, and HLA-B\*38 (AF: 0.174938, 0.128152, 0.080146, 0.064279, and 0.049227), respectively. The first five frequently observed HLA-C alleles were HLA-C\*07, HLA-C\*04, HLA-C\*12, HLA-C\*15, and HLA-C\*06 (AF: 0.230392, 0.137254, 0.112745, 0.102941, and 0.083333), respectively.

The highest frequency was HLA-DRB1\*11 (AF: 0.229569). Subsequently, HLA-DRB1\*04 (AF: 0.147311), HLA-DRB1\*15 (AF: 0.110215), HLA-DRB1\*03 (AF: 0.095698), and HLA-DRB1\*13 (AF: 0.093010). The most common first five HLA-DQA1 alleles were HLA-DQA1\*01, HLA-DQA1\*05, HLA-DQA1\*03, HLA-DQA1\*02, and HLA-DQA1\*04 (AF: 0.365030, 0.325153, 0.193251, 0.098159, and 0.012269), respectively. The most frequently observed HLA-DQB1 allele was HLA-DQB1\*03 (AF: 0.458149), followed by HLA-DQB1\*05 (AF: 0.202643), -DQB1\*06 (AF: 0.178414), -DQB1\*02 (AF: 0.145374), and -DQB1\*04 (AF: 0.015418). We found the same ratios using the R program (Table 1).

We analyzed the frequencies of 336 three-locus haplotypes. Our population was in HWE for all loci using the two methods (Table 2). The haplotype frequencies were similar between the two methods. The most prevalent haplotype

in our population was A\*03 B\*44 DRB1\*04 had the highest frequency (Table 3).

The LD analysis was performed using only the Arlequin software program. According to our LD analysis, there was a significantly strong association between A-B, A-DRB1, and B-DRB1 loci ( $p < 0.001$ ) (Table 4).

## DISCUSSION

In this study, we evaluated HLA allele and haplotype frequencies in the Aegean Region of the Turkish population. This is the first study conducted in the Aegean Region of Türkiye and the first to evaluate the HLA allele and haplotype frequencies of a large Turkish population. In addition, we also compared the two statistical methods for HLA allele and haplotype analyses.

In our study population, the most common HLA-A allele frequency was A\*02, followed by A\*24. The top two alleles were the same in various Turkish populations, including Greece, Bulgaria, Georgia, and in Iran.<sup>14-21</sup> However, only the top allele was the same in the Armenian population.<sup>22</sup> The compatibility of the population in the Aegean Region with surrounding countries may be attributed to past migration. We found that the two most frequently observed HLA-B alleles were B\*35 and B\*51. The HLA-B alleles with the highest frequencies (AF: 0.18) were B\*05 and B\*35 in the Iranian population.<sup>3</sup>

The outcomes were comparable for different Turkish populations, Armenia, Georgia, Iran, and Bulgaria.<sup>15-17,19-22</sup> In the Greek and Turkish populations, the most common allele was HLA-B\*51, followed by B\*35.<sup>17,18</sup> This discrepancy may be attributable to their geographic location. While the most frequent alleles may differ, the predominant B alleles are consistent among populations in the Greek and Aegean Regions. The migration of individuals from Greece and Bulgaria to Türkiye, particularly to the Aegean Region, has played a significant role in fostering diversity throughout history.

The most common HLA-C alleles were HLA-C\*07 and HLA-C\*04, respectively, in our study cohort. However, the results were contrary in the Syrian and Greek populations.<sup>3,18</sup> The most common two alleles were HLA-C\*12 and C\*07 in the Iranian population.<sup>21</sup> The difference in the most common allele may be due to the location of Iran in the Eastern Anatolia Region of Türkiye because our study cohort included the Aegean Region of Türkiye. Despite the replacement of the two most prevalent C alleles in the Greek and Syrian populations, these alleles remained identical to those found in the nations from which the immigrants originated. Due to the significant migration from Syria to the Aegean Region in recent times, the allele frequencies in the population have become comparable. HLA-

**Table 1. Allele frequencies of HLA-A, -B, and -DRB1 by R package degener and Arlequin v.3.5 software**

HLA-A allele	AF	n	HLA-B allele	AF	n	HLA-C allele	AF	n	HLA-DRB1 allele	AF	n	HLA-DQA1 allele	AF	n	HLA-DQB1 allele	AF	n
A*01	0.119885	293	B*07	0.04882	120	C*01	0.029412	6	DRB1*01	0.063441	118	DQA1*01	0.365031	119	DQB1*02	0.145374	66
A*02	0.195990	479	B*08	0.045566	112	C*02	0.063725	13	DRB1*03	0.095699	178	DQA1*02	0.09816	32	DQB1*03	0.45815	208
A*03	0.126432	309	B*13	0.034988	86	C*03	0.058824	12	DRB1*04	0.147312	274	DQA1*03	0.193252	63	DQB1*04	0.015419	7
A*11	0.087561	214	B*14	0.019121	47	C*04	0.137255	28	DRB1*07	0.082796	154	DQA1*04	0.01227	4	DQB1*05	0.202643	92
A*23	0.03477	85	B*15	0.036208	89	C*05	0.029412	6	DRB1*08	0.018817	35	DQA1*05	0.325153	106	DQB1*06	0.178414	81
A*24	0.147708	361	B*18	0.06428	158	C*06	0.083333	17	DRB1*09	0.008065	15	DQA1*06	0.006135	2	In total		454
A*25	0.004909	12	B*27	0.025224	62	C*07	0.230392	47	DRB1*10	0.020968	39	In total		326			
A*26	0.063829	156	B*35	0.174939	430	C*08	0.02451	5	DRB1*11	0.22957	427						
A*29	0.029869	73	B*37	0.013426	33	C*12	0.112745	23	DRB1*12	0.012903	24						
A*30	0.038052	93	B*38	0.049227	121	C*14	0.02451	5	DRB1*13	0.093011	173						
A*31	0.018412	45	B*39	0.013832	34	C*15	0.102941	21	DRB1*14	0.066129	123						
A*32	0.045008	110	B*40	0.032954	81	C*16	0.068627	14	DRB1*15	0.110215	205						
A*33	0.025777	63	B*41	0.019528	48	C*17	0.029412	6	DRB1*16	0.051075	95						
A*34	0.002454	6	B*42	0.000407	1	C*18	0.004902	1	In total		1860						
A*66	0.003682	9	B*44	0.080146	197	In total		204									
A*68	0.041734	102	B*45	0.002848	7												
A*69	0.011047	27	B*46	0.004068	10												
A*74	0.002454	6	B*47	0.002034	5												
A*80	0.000409	1	B*48	0.002848	7												
In total		2444	B*49	0.038649	95												
			B*50	0.031326	77												
			B*51	0.128153	315												
			B*52	0.037022	91												
			B*53	0.009764	24												
			B*54	0.002034	5												
			B*55	0.036208	89												
			B*56	0.002848	7												
			B*57	0.021562	53												
			B*58	0.019935	49												
			B*73	0.002034	5												
			In total		2458												

HLA: Human leukocyte anti-gen, AF: Allele frequency, n: Number of samples

**Table 2. Data table of Hardy-Weinberg equilibrium analysis using the two statistical tools**

Locus	Genotypes	Obs heterozygosity	Exp heterozygosity	p value	SD
By Arlequin v.3.5.					
HLA-A	336	0.88657	0.88908	0.14043	0.00028
HLA-B	336	0.90149	0.92466	0.13648	0.00020
HLA-DRB1	336	0.85672	0.88705	0.33734	0.00054
The R tool					
HLA-A	336	0.88922	0.88782	0.1568	0.017268
HLA-B	336	0.90419	0.92290	0.1622	0.014596
HLA-DRB1	336	0.85928	0.88536	0.4	0.01732
Genotypes: Number of genotypes, Obs: Observed, Exp: Expected, p value: p value for the deviation of HWE, SD: Standard deviation, HLA: Human leukocyte anti-gen					

**Table 3. HLA three-locus haplotype frequency analysis using R and Arlequin tools**

Haplotype	n	Frequency
A*03 B*44 DRB1*04	11	0.016467
A*11, B*35, DRB1*14 A*02 B*51 DRB1*11	10	0.01497
A*33 B*14 DRB1*01 A*23 B*49 DRB1*11	9	0.013473
A*24 B*35 DRB1*11	8	0.011976
A*26 B*38 DRB1*13	7	0.010479
n: Number of samples, HF: Haplotype frequency, HLA: Human leukocyte anti-gen		

**Table 4. Three-loci Linkage disequilibrium estimation by Arlequin software**

Haplotype	D'	p value	Chi-square test values	DF
A-B	0.99990	<0.0001	616.5	432
A-DRB1	0.99990	0.00050	263	192
B-DRB1	0.99921	<0.0001	612.6	324
p<0.05 values were accepted as significant. D': Perfect linkage disequilibrium values, DF: Degrees of freedom				

DRB1\*11 and DRB1\*04 had the highest frequencies in our study group, respectively. The results were in accordance with previous Turkish population studies and in Syrian population.<sup>15-17,23</sup> However, the only highest frequency was the same in border countries as observed in DRB1\*11 and DRB1\*15 in Armenian and Iranian populations, in DRB1\*11 and DRB1\*13 in Georgian populations, and in DRB1\*11 and DRB1\*16 in Greek populations.<sup>18,20-22</sup> It seems that at least one DRB1 allele is identical to those found in nearby nations. This is contingent on the historical migration patterns of adjacent nations. Among the HLA-DQA1 alleles, our results were similar in the Iranian population.<sup>21</sup> However, we could not compare our results with other studies on border populations because the DQA1 locus was not investigated in these studies. The most frequently observed HLA-DQB1 alleles were similar in Iranian, Georgian, Greek, and in other Turkish population studies.<sup>17,18,20,21</sup>

We also evaluated the three-locus haplotype frequencies of ABDRB1 in our population. The most common haplotype differed from Iranian, Armenian, and Georgian populations.<sup>20-22</sup> Our most prevalent haplotype frequency was also different from another Turkish population in Central Anatolia.<sup>24</sup> It is unsurprising that it is identical to the Greek population. These populations have coexisted for ages and have mutually contributed to genetic variety. HLA-A\*02, B\*35, C\*07, DRB1\*03, and DQB1\*03 were the most common alleles in various Caucasian populations.<sup>25-28</sup> Accordingly, this population may have an ancestral relationship with Caucasians. When we considered distant populations, the HLA allele frequencies were related to the Lebanese, Vietnamese, Argentinian, Kazakhstani Russian, Maltese, and Korean populations.<sup>29-34</sup> The similarity of our HLA allele and haplotype frequencies with other populations can be attributed to historical migrations,

genetic admixtures, shared environmental stresses, and evolutionary mechanisms. The balancing selection mechanism and geographical proximity were the primary factors that contributed to these similar characteristics.

The correlation between HLA allele and haplotype frequencies and illness risk is intricate and diverse, as demonstrated by numerous studies. The HLA area is essential for immunological response, with certain alleles potentially enhancing disease susceptibility or providing protection. The interaction is emphasized by the detection of many connections between HLA variants and various autoimmune and infectious illnesses. HLA haplotypes exhibit pleiotropic correlations with numerous diseases, suggesting that certain alleles can continuously affect the risk of different diseases.<sup>35</sup> For example, HLA-B\*08:01 and DRB1\*03:01 are strongly correlated with Graves' disease, although other alleles such as HLA-B\*07:02 have protective benefits.<sup>36</sup> HLA genes are associated with more than 100 disorders, including multiple sclerosis and type 1 diabetes, in which particular alleles may elevate or diminish disease risk. Recent research indicates that multiple sclerosis is linked to DRB1 and DQB1 alleles, which are prevalent in our country.<sup>37</sup> A further study conducted in Russia revealed that the prevalent HLA-C, -DRB1, and -DQB1 alleles in the country were correlated with rheumatoid arthritis, type 1 diabetes, and psoriasis.<sup>38</sup>

The small variations between the two methods for HWE analysis may be due to rounding during the calculation stage. However, these variations are minimal and have no impact on the p value's significance. Both R and Arlequin are capable of performing HLA allele and haplotype frequency, HW, and LD analyses, but they cater to different user needs and expertise levels. Arlequin is a specialized population genetics tool with user-friendly interface. This makes it accessible to researchers with limited programming experience because analyses can be performed with minimal technical input. In contrast, R is a versatile statistical platform that requires additional packages (e.g., genetics, haplo.stats, or LDheatmap). Although this approach offers significant flexibility and allows for the customization of analysis pipelines, it requires advanced programming skills and familiarity with statistical methods. For researchers who are not proficient in R, the learning curve is steep, potentially limiting its immediate usability for LD analysis. Despite these challenges, R's open-source nature and ability to integrate diverse statistical functions makes it an attractive option for experienced users seeking a tailored analytical workflow. By contrast, Arlequin provides an out-of-the-box solution for researchers focusing on LD and other population genetics metrics, particularly when ease of use is a priority.

## Study Limitations

Although our study includes a large cohort of BM donors, it may not fully represent the entire genetic diversity of the Turkish population, as the donors were primarily selected from a specific registry. While we analyzed multiple HLA loci, some additional loci (e.g., HLA-DP) were not included, which could provide further insights into genetic associations and disease susceptibility. Although we compared our results with populations from bordering countries, differences in sampling methodologies and HLA typing techniques could influence the comparability of the findings.

## CONCLUSION

In conclusion, we found that, as expected, the frequencies of the HLA alleles were comparable between the bordering countries. Despite the similarities in allele frequencies, the haplotype frequencies were dissimilar. Knowing a population's allele and haplotype frequencies is crucial for understanding its relationships to transplantation, disease resistance, and therapeutic development. Our findings may assist clinicians in diagnosing disorders, planning transplantation, and creating new medications. Furthermore, numerous statistics software tools are employed to determine allele and haplotype frequencies. Our results show the use of the two Arlequin and R statistical programs to analyze HLA diversity.

## Ethics

**Ethics Committee Approval:** The study was conducted with the permission of İzmir Katip Çelebi University Non-interventional Clinical Research Ethics Committee approved the study (decision no: 0097, date: 23.03.2023).

**Informed Consent:** Informed consent was provided by all patients.

## Footnotes

### Authorship Contributions

Concept: A.Ö.K., T.K.A., İ.P., Design: T.K.A., R.D.G., İ.P., Data Collection or Processing: A.Ö.K., M.S., B.Ç.A., R.D.G., D.İ.T., Analysis or Interpretation: A.Ö.K., T.K.A., M.S., D.İ.T., Literature Search: A.Ö.K., T.K.A., M.S., B.Ç.A., Writing: A.Ö.K.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study received no financial support.

## REFERENCES

1. Satapornpong P, Jinda P, Jantararoungtong T, et al. Genetic diversity of HLA class I and class II alleles in Thai populations:

- contribution to genotype-guided therapeutics. *Front Pharmacol.* 2020;11:78.
2. Varzi AM, Shahsavari F, Tarrahi MJ. Distribution of HLA-DRB1 and HLA-DQB1 alleles in Lak population of Iran. *Hum Immunol.* 2016;77:580-3.
  3. Ikhtiar AM, Jazairi B, Khansa I, Othman A. HLA class I alleles frequencies in the Syrian population. *BMC Res Notes.* 2018;11:324.
  4. Crux NB, Elahi S. Human leukocyte antigen (HLA) and immune regulation: how do classical and non-classical HLA alleles modulate immune response to human immunodeficiency virus and hepatitis C virus infections? *Front Immunol.* 2017;8:832.
  5. Patel JS, Patel MM, Koringa PG, et al. Human leukocyte antigen alleles, genotypes and haplotypes frequencies in renal transplant donors and recipients from West Central India. *Indian J Hum Genet.* 2013;19:219-32.
  6. Patiroğlu T, Akar HH. The frequency of HLA-A, HLA-B, and HLA-DRB1 alleles in patients with acute lymphoblastic leukemia in the Turkish population: a case-control study. *Turk J Hematol.* 2016;33:339-45.
  7. Neamatzadeh H, Dastgheib SA, Mazaheri M, et al. Hardy-Weinberg equilibrium in meta-analysis studies and large-scale genomic sequencing era. *Asian Pac J Cancer Prev.* 2024;25:2229-35.
  8. Abramovs N, Brass A, Tassabehji M. Hardy-Weinberg equilibrium in the large scale genomic sequencing era. *Front Genet.* 2020;11:210.
  9. Hui TYJ, Burt A. Estimating linkage disequilibrium from genotypes under Hardy-Weinberg equilibrium. *BMC Genet.* 2020;21:21.
  10. Anonymous. R: The R project for statistical computing. n.d. Last Accessed Date: 11/22/2024. Available from: <https://www.r-project.org/>
  11. Jombart T. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics.* 2008;24:1403-5.
  12. Paradis E. pegas: an R package for population genetics with an integrated-modular approach. *Bioinformatics.* 2010;26:419-20.
  13. Arrieta-Bolaños E, Hernández-Zaragoza DI, Barquera R. An HLA map of the world: a comparison of HLA frequencies in 200 worldwide populations reveals diverse patterns for class I and class II. *Front Genet.* 2023;14:866407.
  14. Arnaiz-Villena A, Karin M, Bendikuzė N, et al. HLA alleles and haplotypes in the Turkish population: relatedness to Kurds, Armenians and other Mediterraneans. *Tissue Antigens.* 2001;57:308-17.
  15. Kayhan B, Kurtoglu EL, Taskapan H, et al. HLA-A, -B, -DRB1 allele and haplotype frequencies and comparison with blood group antigens in dialysis patients in the East Anatolia region of Turkey. *Transplant Proc.* 2013;45:2123-8.
  16. Uyar FA, Dorak MT, Saruhan-Direskeneli G. Human leukocyte antigen-A, -B and -C alleles and human leukocyte antigen haplotypes in Turkey: relationship to other populations. *Tissue Antigens.* 2004;64:180-7.
  17. Uygun DFK, Özbay M, Sallakçı N, et al. Human leukocyte antigen frequencies in highly sensitized patients. *Turkish J Immunol.* 2015;3:11-4.
  18. Latsoudis H, Stylianakis E, Mavroudi I, et al. Significance of regional population HLA immunogenetic datasets in the efficacy of umbilical cord blood banks and marrow donor registries: a study of Cretan HLA genetic diversity. *Cytherapy.* 2022;24:183-92.
  19. Ivanova M, Spassova P, Michailova A, Naumova E. Distributions of HLA class I alleles and haplotypes in Bulgarians-contribution to understanding the origin of the population. *Tissue Antigens.* 2001;57:208-15.
  20. Rey D, Areces C, Alonso-Rubio J, et al. HLA in Georgians (Caucasus) and their relationship with Eastern Mediterraneans. *Mol Biol Rep.* 2013;40:5523-30.
  21. Abedini F, Rahmanian N, Heidari Z, Feizi A, Sherkat R, Rezaei M. Diversity of HLA class I and class II alleles in Iran populations: systematic review and meta-analysis. *Transpl Immunol.* 2021;69:101472.
  22. Matevosyan L, Chattopadhyay S, Madelian V, et al. HLA-A, HLA-B, and HLA-DRB1 allele distribution in a large Armenian population sample. *Tissue Antigens.* 2011;78:21-30.
  23. Jazairi B, Khansaa I, Ikhtiar A, Murad H. Frequency of HLA-DRB1 and HLA-DQB1 alleles and haplotype association in Syrian population. *Immunol Invest.* 2016;45:172-9.
  24. Yantir E, Gündüz E, Çolak E. HLA alleles, genotype and haplotype analyzes from central Anatolia Region of Turkey. *Balkan Med J.* 2023;40:358-66.
  25. Salvadori L de C, Santana FC de S, Marcos EVC. Frequency of alleles and haplotypes of the human leukocyte antigen system in Bauru, São Paulo, Brazil. *Rev Bras Hematol Hemoter.* 2014;36:108-14.
  26. Cao K, Hollenbach J, Shi X, Shi W, Chopek M, Fernández-Viña MA. Analysis of the frequencies of HLA-A, B, and C alleles and haplotypes in the five major ethnic groups of the United States reveals high levels of diversity in these loci and contrasting distribution patterns in these populations. *Hum Immunol.* 2001;62:1009-30.
  27. Louzada-Junior P, Smith AG, Hansen JA, Donadi EA. HLA-DRB1 and -DQB1 alleles in the Brazilian population of the northeastern region of the state of São Paulo. *Tissue Antigens.* 2001;57:158-62.
  28. Jawdat D, Uyar FA, Alaskar A, Müller CR, Hajeer A. HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 allele and haplotype frequencies of 28,927 Saudi Stem cell donors typed by next-generation sequencing. *Front Immunol.* 2020;11:544768.
  29. Almawi WY, Nemr R, Finan RR, Saldhana FL, Hajje A. HLA-A, -B, -C, -DRB1 and -DQB1 allele and haplotype frequencies in Lebanese and their relatedness to neighboring and distant populations. *BMC Genomics.* 2022;23:456.
  30. Que TN, Khanh NB, Khanh BQ, et al. Allele and haplotype frequencies of HLA-A, -B, -C, and -DRB1 Genes in 3,750 cord blood units from a Kinh Vietnamese population. *Front Immunol.* 2022;13:875283.
  31. Navello M, Müller MC, Riboldi MC, et al. Allele, genotype and haplotype frequencies of the HLA system in the province of Neuquén, Argentina. *Transplantation.* 2022;(95):S566. Available from: <https://cm.tts2022.org/webapp/lecture/1131>
  32. Hajje A, Abdrakhmanova S, Turganbekova A, Almawi WY. HLA allele and haplotype frequencies in Kazakhstani Russians and their relationship with other populations. *HLA.* 2023;101:249-61.
  33. Micallef S, Buttigieg J. #5950 Major HLA allele and haplotype frequencies in the Maltese kidney transplant registry. *Nephrol Dial Transplant.* 2023;(Supplement\_1).
  34. Jung K, Kim JG, Shin S, Roh EY, Hong YJ, Song EY. Allele and haplotype frequencies of 11 HLA loci in Koreans by next-generation sequencing. *HLA.* 2023;101:602-12.

35. Smith CJ, Strausz S; FinnGen; Spence JP, Ollila HM, Pritchard JK. Haplotype analysis reveals pleiotropic disease associations in the HLA region. medRxiv. 2024;24311183.
36. Zawadzka-Starczewska K, Tymoniuk B, Stasiak B, Lewiński A, Stasiak M. Actual associations between HLA haplotype and graves' disease development. J Clin Med. 2022;11:2492.
37. AL-Khaliq IMA. The relation between human leukocyte antigen and multiple sclerosis. SARJMS. 2024;6:54-5.
38. Kazakova P, Smirnova NV, Mitrofanov SI, et al. Study of HLA allele frequencies associated with autoimmune diseases in the population of the russian federation. SSRN Electronic Journal 2022; Available at SSRN: <https://ssrn.com/abstract=4113797> or <http://dx.doi.org/10.2139/ssrn.411379>.